

B (809800,826000)



500m Study Area

A (809800,824600)



Scale: As shown

Figure 7-K1 Cumulative 1-hour Ni Concentration ( $\mu\text{g}/\text{m}^3$ ) at 8.5mPD emitted from Source Location A4 (Mitigated Scenario 2)



B (809800,826000)



500m Study Area

A (809800,824600)



Scale: As shown  
Figure 7-K2 Cumulative 1-hour Ni Concentration ( $\mu\text{g}/\text{m}^3$ ) at 8.5mPD emitted from Source Location A6 (Mitigated Scenario 2)



B (809800,826000)



500m Study Area

A (809800,824600)



Scale: As shown  
Figure 7-K3 Cumulative 1-hour Ni Concentration ( $\mu\text{g}/\text{m}^3$ ) at 8.5mPD emitted from Source Location A8 (Mitigated Scenario 2)



B (809800,826000)



500m Study Area

A (809800,824600)



Scale: As shown  
Figure 7-K4 Cumulative 1 Hour Ni Concentration ( $\mu\text{g}/\text{m}^3$ ) at 22.5mPD emitted from Source Location A9 (Mitigated Scenario 2)



B (809800,826000)



500m Study Area


A (809800,824600)



Scale: As shown

Figure 7-K5 Cumulative 1-hour Ni Concentration ( $\mu\text{g}/\text{m}^3$ ) at 61.5mPD emitted from Source Location A2 (Mitigated Scenario 2)





B (809800,826000)



500m Study Area

A (809800,824600)



Scale: As shown

Figure 7-K6 Cumulative 1-hour Ni Concentration ( $\mu\text{g}/\text{m}^3$ ) at 61.5mPD emitted from Source Location A5 (Mitigated Scenario 2)