

BCG Vaccination-Reaction and Care

Within two to four weeks after vaccination, a small red papule or ulcer may develop which will gradually dry up during the following few weeks, then heal completely leaving very small or no scar. Occasionally a few children may develop lumps under the armpit caused by enlarged glands. Other adverse reactions are very rare.

The following points should be noted after BCG vaccination:

1. Don't worry in case of minor pustule or ulcer formation. This is a normal reaction which will heal gradually after a few weeks.
2. Bath should be taken as usual. The vaccination site should be kept clean and dry. If necessary, it can be cleansed with distilled water and dried up with gauze afterwards.
3. **No** medication or ointment should be applied to the vaccination site. The site should **not** be covered with adhesive plaster or dressing. Tight clothing should be avoided.

For queries about the aforesaid reactions, please telephone or make a personal visit to the nearby Chest Clinic for further information or examination.

Chest Clinics of Tuberculosis and Chest Service:	Tel. No.
Wanchai Chest Clinic	1/F, Wanchai Polyclinic, 99 Kennedy Road, Wan Chai, H.K. 2591 1173
Sai Ying Pun Chest Clinic	2/F, Sai Ying Pun Jockey Club Polyclinic, 134 Queen's Road West, H.K. 2540 9778
Shaukiwan Chest Clinic	2/F, Shaukiwan Jockey Club Clinic, 8 Chai Wan Road, H.K. 2535 3839
Kowloon Chest Clinic	G/F, 147A Argyle Street, Kln. 2767 5712
Yaumatei Chest Clinic	2/F, Yaumatei Jockey Club Polyclinic, 145 Battery Street, Kln. 2245 9527
Shek Kip Mei Chest Clinic	1/F, Shek Kip Mei Polyclinic, 2 Berwick Street, Service suspension till further announcement 2779 7257
East Kowloon Chest Clinic	G/F, East Kowloon Polyclinic, 160 Hammer Hill Road, Kln. 2325 6851
Yung Fung Shee Chest Clinic	1/F, Yung Fung Shee Memorial Centre, 79 Cha Kwo Ling Road, Kln. 2727 8252
Yan Oi Chest Clinic	2/F, Yan Oi Polyclinic, 6 Tuen Lee Street, Tuen Mun, N.T. 2404 3747
South Kwai Chung Chest Clinic	G/F, 310 Kwai Shing Circuit, South Kwai Chung, N.T. 2420 6049
Yuen Chau Kok Chest Clinic	1/F, 29 Chap Wai Kon Street, Sha Tin, N.T. 2686 7432
Tai Po Chest Clinic	1/F, Tai Po Jockey Club Clinic, 37 Ting Kok Road, Tai Po, N.T. 2689 5601

卡介苗防疫注射的反應與護理

注射卡介苗後二至四星期內，注射部位可能呈現小紅丘疹或潰瘍，此情況出現後，需待數星期才會逐漸退卻及完全痊癒，留下極輕微疤痕或甚至不留疤痕。間中有些兒童會因淋巴腺擴大以致腋窩下腫起幾處。其他不良反應，則極為罕見。

接受卡介苗注射後，請注意：

1. 如有灌膿或小潰瘍情況出現，請勿驚惶，因這是正常現象。數星期後，上述情況便會逐漸痊癒。
2. 可照常沐浴。注射部位應保持清潔乾爽，需要時可用凍開水清潔並以紗布吸乾水份。
3. 在注射部位切勿塗搽任何藥物、藥水及藥膏等，毋需貼上膠布或敷上任何敷料，並避免緊身衣物。

如果對上述反應有任何疑慮，請致電或前往就近的胸肺服務診所查詢或檢查。

胸肺服務診所：

電話：

灣仔胸肺科診所	香港灣仔堅尼地道 99 號灣仔分科診所 1 樓	2591 1173
西營盤胸肺科診所	香港西營盤皇后大道西 134 號西營盤賽馬會分科診所 2 樓	2540 9778
筲箕灣胸肺科診所	香港筲箕灣柴灣道 8 號筲箕灣賽馬會診所 2 樓	2535 3839
九龍胸肺科診所	九龍旺角亞皆老街 147A 地下	2767 5712
油麻地胸肺科診所	九龍油麻地炮台街 145 號油麻地賽馬會分科診所 2 樓	2245 9527
石硤尾胸肺科診所	九龍石硤尾巴域街 2 號 待重建中，暫停所有服務。	2779 7257
東九龍胸肺科診所	九龍鑽石山斧山道 160 號東九龍分科診所地下	2325 6851
容鳳書胸肺科診所	九龍觀塘茶果嶺道 79 號容鳳書紀念中心 1 樓	2727 8252
仁愛胸肺科診所	新界屯門屯利街 6 號仁愛分科診所 2 樓	2404 3747
南葵涌胸肺科診所	新界葵涌葵盛圍 310 號地下	2420 6049
圓洲角胸肺科診所	新界沙田插桅桿街 29 號 1 樓	2686 7432
大埔胸肺科診所	新界大埔汀角路 37 號大埔賽馬會診所 1 樓	2689 5601